

Attachment: Exam questions

Table 1: Module-related exam questions at the end of the semester

Summer semester 2020 (L1 ^a)	Winter semester 2020/21 (L2 & L3 ^a)
<p>[TypeA] Which statement about alcohol consumption is most accurate?</p> <ol style="list-style-type: none"> 1. A standard glass of alcohol contains approx. 20 g of pure alcohol 2. According to the definition of the German Centre for Addiction Issues (Deutsche Hauptstelle für Suchtfragen - DHS), "low-risk consumption" among men means up to 3 standard drinks per drinking session. 3. About 50 litres of pure alcohol are drunk per capita per year in Germany. 4. At-risk consumption affects about 2% of Germans between the ages of 18 and 64. (Correct) 5. A litre glass of beer contains approx. 60 g of pure alcohol. 	<p>[TypeA] Mr Miller (60 years old, in employment) comes to your family practice today. In addition to cigarettes, Mr Miller states that he consumes a bottle of wine almost every day. They suspect dependency. Which of the following questions is <u>not</u> part of the CAGE questionnaire?</p> <ol style="list-style-type: none"> 1. Have you ever had problems at work because of drinking? (Correct) 2. Have you ever felt guilty about drinking? 3. Have you ever felt you needed to cut down on your drinking? 4. Have you ever felt the needed a drink first thing in the morning to become fit and alert? 5. Have people annoyed you by criticising your drinking?
<p>[TypeA] You are a general practitioner in a practice. Ms Smith, a chemist with a PhD and a good income, comes to you. She hopes that today's appointment will help her to stop smoking, as she is unable to stop smoking permanently despite being pregnant (6 weeks). Which statement does not apply?</p> <ol style="list-style-type: none"> 1. This makes her one of 12 % of women in Germany who smoke during pregnancy, regardless of their social status. 2. Irrespective of pregnancy and social status, this places her among the 20 % of women who smoke in Germany. 3. In the transtheoretical model, she is therefore in the preparation stage. 4. She is therefore among only less than 3% of pregnant women with high social status who smoke during pregnancy. 5. You can help her to stop smoking quickly by prescribing a medicine such as varenicline. (Correct) 	<p>[TypeA] The test gives no indication of dependence. According to the amount he drinks, however, Mr. Miller has a high-risk level of consumption, but does not wish to change the amount he drinks in the near future. This places him at the precontemplation stage in Prochaska and di Clemente's model.</p> <p>What is the most important thing you should do at this juncture?</p> <ol style="list-style-type: none"> 1. Above all, point out to him the consequences of his current level of consumption. 2. Persuade him to drink less now. 3. Agree on concrete goals for reducing the amount he drinks. 4. Let him make suggestions on how he can reduce his consumption. 5. Arouse interest in the topic of alcohol consumption by providing information. (Correct)

^a L: Learning objectives according to Table 1.